Please fax both pages of completed form to your team at 866.489.1901.

To reach your team, call toll-free 866.489.1899.

You can now monitor shipments and chat online if you have questions. Go to MyAccredoPatients.com to log in or get started.

Prescription & Enrollment Form

Inflammatory Bowel Disease



Four simple steps to submit your referral.

1 Patient Information			Please provide copies of fro and prescription insurance of	
New patient				
Patient's first name		Last name _		Middle initial
Male Female Last 4 digits of SSN			Date of birth	
Street address				Apt #
City		State		Zip
Home phone	Cell phone		E-mail address	
Parent/guardian (if applicable)				
Home phone	Cell phone		E-mail address	
Alternate caregiver/contact				
Home phone	Cell phone		E-mail address	
OK to leave message with alternate care	egiver/contact			
Patient's primary language: English	Other If other	, please specify		
Date Time				
Prescriber info: Prescriber's first name				
Prescriber's title				
Office phone				
Office contact and title				
Office street address				
City				•
			sion site, complete information	
Infusion info: Infusion site name				
Site street address				
City				
Infusion clinic contact name				•
3 Clinical Information				
Primary ICD-10 code (REQUIRED):		Rationale	e for therapy	
Has the patient been treated previously fo	r this condition?	Yes No Is pa	tient currently on therapy?	Yes No
Please list all therapies tried/failed:				
Patient wt Date wt				BSAm2
NKDA Known drug allergies				
Concurrent meds				

Patient's first name Las	ast name	Middle initial	Date of birth
Prescriber's first name	Last name	Phone	9

4

Prescribing Information

Medication	Dose	Directions	Quantity/Refills
Simponi® (golimumab)	100mg	Initial Dose: 200mg subcutaneously at week 0 followed by 100mg subcutaneously at 2 weeks Maintenance Dose: 100mg subcutaneously every 4 weeks	1-month supply 3-month supply Other Refills
Xeljanz® (tofacitinib citrate)	5mg tablets 10mg tablets	Take 10mg by mouth twice daily for 8 weeks, followed by 5mg twice daily Take 10mg by mouth twice daily Take 5mg by mouth twice daily Take 5mg by mouth once daily	1-month supply 3-month supply Other Refills
Other			
Ancillary Supplies: (Prescriber to strike through if not required) Dispense ancillary supplies such as needles, syringes, sterile water, etc. and home medical equipment necessary to administer the therapy as needed.		Send quantity sufficient for medication days supply.	

If shipped to physician's office, physician accepts on behalf of patient for administration in office.

Prescriber's signature required (sign below) (Physician attests this is his/her legal signature. NO STAMPS)

SIGN				
HERE	Date	Dispense as written	Date	Substitution allowed

The prescriber is to comply with his/her state-specific prescription requirements such as e-prescribing, state-specific prescription form, fax language, etc. Non-compliance with state-specific requirements could result in outreach to the prescriber.

