

# Referral Form for TYVASO (treprostinil) and TYVASO DPI (treprostinil)



Tyvaso and Tyvaso DPI are available only through select Specialty Pharmacy Services (SPS) providers. Follow these 5 steps to complete each section of the following referral form.



## GET STARTED CHECKLIST

- 1** Fill out the Patient and Insurance Information. Let your patient know that an SPS provider will be calling and it is important to answer or return the call.
- 2** Complete and sign the Prescriber Information, Medical Information, and Treatment History and Transition Statement.
- 3** Complete and sign the Prescription Information, Statement of Medical Necessity for either **PH-ILD** or **PAH**, and Calcium Channel Blocker Statement (CCB Statement not required for PH-ILD).
- 4** Complete the Optional Side Effect Management page.
- 5** Attach the clinical documents outlined on the **Fax Cover Sheet**, including right heart catheterization test results, history and physical, and echocardiogram results. Use the included **Fax Cover Sheet** in this PDF to fax the referral form and signed supporting documents to your preferred SPS provider. (Insurance plans vary and may impact the approval process.)

## STEP 1 PATIENT INFORMATION

Name - First			Middle	Last	
Date of Birth			Gender	Last 4 Digits of SSN	
Home Address					
City		State		Zip	
Shipping Address (if different from home address)					
City		State		Zip	
Telephone: <input type="checkbox"/> Home <input type="checkbox"/> Cell <input type="checkbox"/> Work		Alternate Telephone: <input type="checkbox"/> Home <input type="checkbox"/> Cell <input type="checkbox"/> Work		Best Time to Call: <input type="checkbox"/> Morning <input type="checkbox"/> Afternoon <input type="checkbox"/> Evening	
E-mail Address				Okay to leave a voicemail? <input type="checkbox"/> Yes <input type="checkbox"/> No	
Caregiver/Family Member		Caregiver Telephone: <input type="checkbox"/> Home <input type="checkbox"/> Cell <input type="checkbox"/> Work		Caregiver Alternate Telephone: <input type="checkbox"/> Home <input type="checkbox"/> Cell <input type="checkbox"/> Work	
Caregiver E-mail Address		Caregiver Alternate E-mail Address		Okay to leave a voicemail? <input type="checkbox"/> Yes <input type="checkbox"/> No	

## STEP 1 INSURANCE INFORMATION

Primary Prescription Insurance					
Subscriber ID #			Group #	Telephone	
Primary Medical Insurance			Policy Holder/Relationship		
Subscriber ID #			Group #	Telephone	
Secondary Medical Insurance			Policy Holder/Relationship		
Subscriber ID #			Group #	Telephone	

Please include copies of the front and back of the patient's medical and prescription insurance card(s).



Patient Name: \_\_\_\_\_ Date of Birth: \_\_\_\_\_

**STEP 3 PH-ILD - USE THIS SECTION FOR PH-ILD**

**Diagnosis - The following ICD-10 codes do not suggest approval, coverage, or reimbursement for specific uses or indications.**

Please include one PH-specific diagnosis code **AND** one ILD-specific diagnosis code.

**PH Diagnosis Codes:**

ICD-10 I27.23  Pulmonary hypertension due to lung diseases and hypoxia Other ICD-10: \_\_\_\_\_

**ILD Diagnosis Codes:**

**IIP:** ICD-10 J84.10  Pulmonary fibrosis, unspecified ICD-10 J84.111  Idiopathic interstitial pneumonia, NOS ICD-10 J84.112  Idiopathic pulmonary fibrosis

**CTD-related ILD:** ICD-10 M34.81  Systemic sclerosis with lung involvement

**Environmental/Occupational Lung Disease:**

ICD-10 J61  Pneumoconiosis due to asbestos and other mineral fibers ICD-10 J67.9  Hypersensitivity pneumonitis due to unspecified dust

**Other Causes:** ICD-10 J17  Pneumonia in disease classified elsewhere Other ICD-10: \_\_\_\_\_

Please visit [www.utassist.com/codes](http://www.utassist.com/codes) for additional ICD-10 codes related to PAH, PH, and ILD



**TYVASO (treprostinil) 1.74mg/2.9ml ampule (0.6mg/ml) Inhalation Solution**

**Target dose: 9 breaths (54 mcg) to 12 breaths (72 mcg), 4 times daily** - Start with 3 breaths (18 mcg) 4 times daily (if 3 breaths are not tolerated, use 1 to 2 breaths). Increase by an additional 1 breath per week, as tolerated, until the target dose of 9 breaths (54 mcg) to 12 breaths (72 mcg), 4 times daily is achieved.

- TYVASO Inhalation System Starter Kit (28-day supply) 0 refills
- TYVASO Inhalation System Refill Kit (28-day supply) X \_\_\_\_\_ refills

Prescriber may specify any alternative or additional dosing and titration instructions here:

**OR**

**TYVASO DPI (treprostinil) Inhalation Powder**

**Target dose:**  48 mcg or  64 mcg or  Other \_\_\_\_\_ mcg per treatment session, 4 times daily (Check One)

Start with one 16-mcg cartridge per treatment session, 4 times daily. Increase cartridge strength by 16 mcg per treatment session every 1 to 2 weeks, as tolerated, to selected target dose. Titration schedule may vary based on tolerability. If the prescribed dose is higher than 64 mcg per treatment session, more than 1 cartridge will be needed per session.

**TYVASO DPI Titration Kit (28-day supply) Choose for titration phase.**

- 16 mcg (112 ct), 32 mcg (112 ct), and 48 mcg (28 ct) 1 refill

**TYVASO DPI Maintenance Kit (28-day supply)** X \_\_\_\_\_ refills

**Inhale one breath per cartridge, 4 times daily. Please check the box of the maintenance kit for the desired target dose.**

- 16 mcg (112 ct)  32 mcg (112 ct)  48 mcg (112 ct)  64 mcg (112 ct)

Prescriber may specify any alternative or additional dosing and titration instructions on the line below. If the prescribed dose is higher than 64 mcg per treatment session, more than 1 cartridge will be needed per session:

*Specialty Pharmacy to contact prescribing practitioner for adjustments to the written orders specified above.*

**Dose Comparison**

TYVASO Nebulizer # of Breaths	TYVASO DPI Cartridge Strength
≤5	16 mcg
6 to 7	32 mcg
8 to 10	48 mcg
11 to 12	64 mcg

**NURSING ORDERS**

RN visit to provide assessment and education on administration, dosing, and titration. **Location:**  Home  Outpatient Clinic  Hospital

The Prescriber is to comply with their state-specific prescription requirements, such as e-prescribing, state-specific prescription form, fax language, etc. Non-compliance of state-specific requirements could result in outreach to the Prescriber.

CHECK ONE

**Nurse Visits**

Specialty Pharmacy home healthcare RN visit to provide education on self-administration of Tyvaso or Tyvaso DPI, including dose, titration, and side effect management.

**OR**

Prescriber directed Specialty Pharmacy home healthcare RN visit(s) as detailed below:

\_\_\_\_\_

**STEP 3 PRESCRIBER SIGNATURE: PRESCRIPTION AND STATEMENT OF MEDICAL NECESSITY**

SIGN HERE

I certify that the pulmonary hypertension associated with interstitial lung disease therapy ordered above is medically necessary and that I am personally supervising the care of this patient. **PHYSICIAN'S SIGNATURE REQUIRED TO VALIDATE PRESCRIPTIONS.**

Physician's Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Dispense as Written

Substitution Allowed

DAW

State-Specific Dispense as Written (DAW) Selection Verbiage: \_\_\_\_\_

(Physician attests this is his/her legal signature. NO STAMPS.) PRESCRIPTIONS MUST BE FAXED.

*Tyvaso and Tyvaso DPI are registered trademarks of United Therapeutics Corporation. All other brands are trademarks of their respective owners. The makers of these brands are not affiliated with and do not endorse United Therapeutics or its products.*

Patient Name: \_\_\_\_\_ Date of Birth: \_\_\_\_\_

**STEP 3 PAH - USE THIS SECTION FOR PAH**

Diagnosis - The following ICD-10 codes do not suggest approval, coverage, or reimbursement for specific uses or indications.

- ICD-10 I27.0 Primary pulmonary hypertension:  Idiopathic PAH  Heritable PAH  
 ICD-10 I27.21 Secondary pulmonary arterial hypertension:  Connective tissue disease  Drugs/Toxins induced  Portal hypertension  HIV  Congenital heart diseases  
 Other: \_\_\_\_\_ Other ICD-10: \_\_\_\_\_

Please visit [www.utassist.com/codes](http://www.utassist.com/codes) for additional ICD-10 codes related to PAH, PH, and ILD



**TYVASO (treprostinil) 1.74mg/2.9ml ampule (0.6mg/ml) Inhalation Solution**

**Target dose: 9 breaths (54 mcg) to 12 breaths (72 mcg), 4 times daily** - Start with 3 breaths (18 mcg) 4 times daily (if 3 breaths are not tolerated, use 1 to 2 breaths). Increase by an additional 3 breaths every week, if tolerated, until the target dose of 9 breaths (54 mcg) to 12 breaths (72 mcg), 4 times daily.

- TYVASO Inhalation System Starter Kit (28-day supply) 0 refills  
 TYVASO Inhalation System Refill Kit (28-day supply) X \_\_\_\_\_ refills

Prescriber may specify any alternative or additional dosing and titration instructions here:

**OR**



**TYVASO DPI (treprostinil) Inhalation Powder**

**Target dose:  48 mcg or  64 mcg or  Other \_\_\_\_\_ mcg per treatment session, 4 times daily (Check One)**  
 Start with one 16-mcg cartridge per treatment session, 4 times daily. Increase cartridge strength by 16 mcg per treatment session every week to selected target dose. Titration schedule may vary based on tolerability. If the prescribed dose is higher than 64 mcg per treatment session, more than 1 cartridge will be needed per session.

**TYVASO DPI Titration Kit (28-day supply) Choose for titration phase.**

- 16 mcg (112 ct), 32 mcg (112 ct), and 48 mcg (28 ct) 1 refill

**TYVASO DPI Maintenance Kit (28-day supply) X \_\_\_\_\_ refills**

**Inhale one breath per cartridge, 4 times daily. Please check the box of the maintenance kit for the desired target dose.**

- 16 mcg (112 ct)  32 mcg (112 ct)  48 mcg (112 ct)  64 mcg (112 ct)

Prescriber may specify any alternative or additional dosing and titration instructions on the line below. If the prescribed dose is higher than 64 mcg per treatment session, more than 1 cartridge will be needed per session:

Specialty Pharmacy to contact prescribing practitioner for adjustments to the written orders specified above.

**Dose Comparison**

TYVASO Nebulizer # of Breaths	TYVASO DPI Cartridge Strength
≤5	16 mcg
6 to 7	32 mcg
8 to 10	48 mcg
11 to 12	64 mcg

**NURSING ORDERS**

RN visit to provide assessment and education on administration, dosing, and titration. **Location:**  Home  Outpatient Clinic  Hospital

The Prescriber is to comply with their state-specific prescription requirements, such as e-prescribing, state-specific prescription form, fax language, etc. Non-compliance of state-specific requirements could result in outreach to the Prescriber.

CHECK ONE

**Nurse Visits**

Specialty Pharmacy home healthcare RN visit to provide education on self-administration of Tyvaso or Tyvaso DPI, including dose, titration, and side effect management.

**OR**  Prescriber directed Specialty Pharmacy home healthcare RN visit(s) as detailed below:

**STEP 3 CALCIUM CHANNEL BLOCKER STATEMENT (Not required for PH-ILD patients)**

Please indicate below if the Patient named above was trialed on a Calcium Channel Blocker prior to the initiation of therapy and indicate the results.

**A Calcium Channel Blocker was not trialed because:**

- Patient has depressed cardiac output  Patient is hemodynamically unstable or has a history of postural hypotension  
 Patient has systemic hypotension  Patient did not meet ACCP Guidelines for Vasodilator Response  
 Patient has known hypersensitivity  Patient has documented bradycardia or second- or third-degree heart block  
 Other: \_\_\_\_\_

**OR**

**The following Calcium Channel Blocker was trialed:** \_\_\_\_\_

With the following response(s):

- Patient hypersensitive or allergic \_\_\_\_\_  Pulmonary arterial pressure continued to rise  
 Adverse event \_\_\_\_\_  Patient became hemodynamically unstable  
 Disease continued to progress or patient remained symptomatic \_\_\_\_\_  
 Other: \_\_\_\_\_

**STEP 3 PRESCRIBER SIGNATURE: PRESCRIPTION AND STATEMENT OF MEDICAL NECESSITY**

I certify that the pulmonary arterial hypertension therapy ordered above is medically necessary and that I am personally supervising the care of this patient.

**PHYSICIAN'S SIGNATURE REQUIRED TO VALIDATE PRESCRIPTIONS.**

Physician's Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Dispense as Written

Substitution Allowed

SIGN HERE

DAW

State-Specific Dispense as Written (DAW) Selection Verbiage: \_\_\_\_\_

(Physician attests this is his/her legal signature. NO STAMPS.) PRESCRIPTIONS MUST BE FAXED.

Tyvaso and Tyvaso DPI are registered trademarks of United Therapeutics Corporation. All other brands are trademarks of their respective owners. The makers of these brands are not affiliated with and do not endorse United Therapeutics or its products.



Fax the completed referral form and documentation to the Specialty Pharmacy of your choice below.

**STEP 5 FAX COVER SHEET**

**Date:** \_\_\_\_\_ **Patient Initials:** \_\_\_\_\_ **Patient Date of Birth:** \_\_\_\_\_

**To: (check one)**       **Accredo Health Group, Inc.**       **CVS Specialty**  
 Fax: 1-800-711-3526      Fax: 1-877-943-1000  
 Phone: 1-866-344-4874      Phone: 1-877-242-2738

**From:** (Name of agent of prescriber who transmitted the facsimile/prescription)  
 \_\_\_\_\_

**Facility Name:** \_\_\_\_\_

**Fax:** \_\_\_\_\_

**Included in this fax:**

- Completed Tyvaso and Tyvaso DPI Therapy Referral Form including**
  - Step 1 - Patient Information and Insurance Information (including front and back copies of medical and prescription insurance card(s))
  - Step 2 - Prescriber Information, Medical Information/Patient Evaluation/Supporting Documentation, and Treatment History and Transition Statement
  - Step 3 - Prescription Information and Calcium Channel Blocker Statement (CCB Statement not required for PH-ILD)
  - Step 4 - Optional Side Effect Management
  
- Included signed and dated documents**
  - Right Heart Catheterization Results
  - History and Physical (including Onset of Symptoms, PAH or PH associated with ILD Clinical Signs and Symptoms, Course of Illness)
  - Need for Specific Drug Therapy and 6-minute walk test results (6-minute walk test not required for PH-ILD)
  - Echocardiogram Results (not required for PH-ILD patients)
  - High-Resolution CT Scan (not required for PAH patients)

**Number of Pages:** \_\_\_\_\_

**Additional Comments:**  
 \_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_

US-TYV-0710