

Please fax both pages of completed form to your team at 888.302.1028.

To reach your team, call toll-free 844.516.3319.

You can now monitor shipments and chat online if you have questions. Go to [MyAccredoPatients.com](https://MyAccredoPatients.com) to log in or get started.

## Prescription & Enrollment Form Ulcerative Colitis

accredo®

Four simple steps to submit your referral.

### 1 Patient Information



Please provide copies of front and back of all medical and prescription insurance cards.

New patient      Current patient

Patient's first name \_\_\_\_\_ Last name \_\_\_\_\_ Middle initial \_\_\_\_\_

Preferred patient first name \_\_\_\_\_ Preferred patient last name \_\_\_\_\_

Sex at birth:    Male    Female    Gender identity \_\_\_\_\_ Pronouns \_\_\_\_\_ Last 4 digits of SSN \_\_\_\_\_

Date of birth \_\_\_\_\_ Street address \_\_\_\_\_ Apt # \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_

Home phone \_\_\_\_\_ Cell phone \_\_\_\_\_ Email address \_\_\_\_\_

Parent/guardian (if applicable) \_\_\_\_\_

Home phone \_\_\_\_\_ Cell phone \_\_\_\_\_ Email address \_\_\_\_\_

Alternate caregiver/contact \_\_\_\_\_

Home phone \_\_\_\_\_ Cell phone \_\_\_\_\_ Email address \_\_\_\_\_

OK to leave message with alternate caregiver/contact

Patient's primary language:    English    Other    If other, please specify \_\_\_\_\_

### 2 Prescriber Information

All fields must be completed to expedite prescription fulfillment.

Date \_\_\_\_\_ Time \_\_\_\_\_ Date medication needed \_\_\_\_\_

Office/clinic/institution name \_\_\_\_\_

**Prescriber info:** Prescriber's first name \_\_\_\_\_ Last name \_\_\_\_\_

Prescriber's title \_\_\_\_\_ If NP or PA, under direction of Dr. \_\_\_\_\_

Office phone \_\_\_\_\_ Fax \_\_\_\_\_ NPI # \_\_\_\_\_ License # \_\_\_\_\_

Office contact and title \_\_\_\_\_ Office contact email \_\_\_\_\_

Office street address \_\_\_\_\_ Suite # \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_

Infusion location:    Patient's home    Prescriber's office    Infusion site    If infusion site, complete information below dotted line:  
-----

**Infusion info:** Infusion site name \_\_\_\_\_ Clinic/hospital affiliation \_\_\_\_\_

Site street address \_\_\_\_\_ Suite # \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_

Infusion site contact \_\_\_\_\_ Phone \_\_\_\_\_ Fax \_\_\_\_\_ Email \_\_\_\_\_

### 3 Clinical Information

**Primary ICD-10 code (REQUIRED):** \_\_\_\_\_ Has the patient been treated previously for this condition?    Yes    No

Is patient currently on therapy?    Yes    No    Please list all therapies tried/failed: \_\_\_\_\_

Patient wt \_\_\_\_\_ Date wt obtained \_\_\_\_\_

NKDA    Known drug allergies \_\_\_\_\_

Concurrent meds \_\_\_\_\_

Patient's first name \_\_\_\_\_ Last name \_\_\_\_\_ Middle initial \_\_\_\_\_ Date of birth \_\_\_\_\_

Prescriber's first name \_\_\_\_\_ Last name \_\_\_\_\_ Phone \_\_\_\_\_

## 4 Prescribing Information

Medication	Strength/Formulation	Directions	Quantity/Refills
Simponi® (golimumab)	100mg/mL in each single-dose prefilled syringe (PFS) 100mg/mL in each single-dose pen	<b>Loading dose:</b> Inject 200mg subcutaneously at week 0, followed by 100mg subcutaneously at week 2	QS for 42-day supply loading dose No Refills
		<b>Maintenance dose:</b> Inject 100mg subcutaneously every 4 weeks.	1-month supply 3-month supply Refill QS 1 year unless otherwise noted Other _____
Stelara® (ustekinumab)	90mg/mL in each single-dose PFS	<b>Maintenance dose:</b> Inject 90mg subcutaneously every 8 weeks.	1-month supply 3-month supply Refill QS 1 year unless otherwise noted Other _____
		<b>Maintenance Dose Only Needed.</b> If loading dose is needed, please see IV referral form. By selecting Stelara on this form, I am indicating that patient has already received/does not need IV loading dose at this time.	
Xeljanz®	10mg tablets	<b>Loading dose:</b> Take 10mg by mouth twice daily for 8 weeks, followed by 5mg twice daily	QS for 2-month loading dose No Refills
	5mg tablets 10mg tablets	<b>Maintenance dose:</b> Take 10mg by mouth twice daily Take 5mg by mouth twice daily Take 5mg by mouth once daily	1-month supply 3-month supply Refill QS 1 year unless otherwise noted Other _____
Xeljanz XR™	22mg ER tablets	<b>Loading dose:</b> 22mg once daily for at least 8 weeks, followed by 11mg once daily	QS for 2-month loading dose No Refills
	11mg ER tablets	<b>Maintenance dose:</b> Take 11mg by mouth once daily	1-month supply 3-month supply Refill QS 1 year unless otherwise noted Other _____
Zeposia® (ozanimod)	<b>Starter dose:</b> Starter Pack (28 day) Starter Pack (7 day)	Take one 0.23mg capsule daily for 4 days, then one 0.46mg capsule for 3 days, then one 0.92mg capsule daily thereafter	1 kit No Refills
	<b>Maintenance dose:</b> 0.92mg capsules	Take one capsule daily	1-month supply 3-month supply Refill QS 1 year unless otherwise noted Other _____
Other			

Dispense needles, syringes, ancillary supplies and home medical equipment necessary to administer medication.

If shipped to physician's office, physician accepts on behalf of patient for administration in office.

**Prescriber's signature required (sign below) (Physician attests this is his/her legal signature. NO STAMPS)**SIGN  
HERE

Date

Dispense as written

Date

Substitution allowed

The prescriber is to comply with his/her state-specific prescription requirements such as e-prescribing, state-specific prescription form, fax language, etc. Non-compliance with state-specific requirements could result in outreach to the prescriber.