

Please fax both pages of completed form to your team at 866.531.1025.

To reach your team, call toll-free 866.839.2162.

You can now monitor shipments and chat online if you have questions. Go to MyAccredoPatients.com to log in or get started.

Prescription & Enrollment Form
Xolair® (omalizumab)

accredo®

Four simple steps to submit your referral.

1 Patient Information



Please provide copies of front and back of all medical and prescription insurance cards.

New patient Current patient

Patient's first name _____ Last name _____ Middle initial _____

Sex at birth: Male Female Preferred pronouns _____ Last 4 digits of SSN _____ Date of birth _____

Street address _____ Apt # _____

City _____ State _____ Zip _____

Home phone _____ Cell phone _____ Email address _____

Parent/guardian (if applicable) _____

Home phone _____ Cell phone _____ Email address _____

Alternate caregiver/contact _____

Home phone _____ Cell phone _____ Email address _____

OK to leave message with alternate caregiver/contact

Patient's primary language: English Other If other, please specify _____

2 Prescriber Information

All fields must be completed to expedite prescription fulfillment.

Date _____ Time _____ Date medication needed _____

Office/clinic/institution name _____

Prescriber's first name _____ Last name _____

Prescriber's title _____ If NP or PA, under direction of Dr. _____

Office phone _____ Fax _____ NPI # _____ License # _____

Office contact and title _____ Office contact email _____

Office street address _____ Suite # _____

City _____ State _____ Zip _____

Deliver product to: Prescriber's office Patient's home

3 Clinical Information

ICD-10 code (REQUIRED): _____

NKDA Known drug allergies _____

Prior anaphylactic reaction: No Yes (Reason/date _____)

Concurrent meds _____

Concomitant therapies: Short-acting beta agonist Long-acting beta agonist Antihistamines Decongestants Immunotherapy

Inhaled corticosteroid Leukotriene modifiers Oral steroids Nasal steroids Other _____

Lab results: History of positive skin OR RAST test to a perennial aeroallergen

Pre-treatment serum IgE level _____ IU per mL Test date _____ Pre-treatment serum eosinophils _____ cells/mL

and/or sputum eosinophils _____ Date _____ Patient wt _____ kg Date wt obtained _____

MD Specialty (required): Allergist Pulmonologist ENT Primary care Pediatrician Other _____

Prescription type: Naïve/new start Restart Continued therapy

Patient's first name _____ Last name _____ Middle initial _____ Date of birth _____

Prescriber's first name _____ Last name _____ Phone _____

4 Prescribing Information

Medication	Strength/Formulation	Directions	Quantity/Refills
Xolair® (omalizumab)	Prefilled syringe <i>Pharmacy to dispense the least amount of syringes to complete total dose. Prefilled syringe available in 75mg and 150mg.</i> 150mg single dose vial	Every 4 weeks dosing: Inject 75mg per dose under the skin every 4 weeks Inject 150mg per dose under the skin every 4 weeks Inject 225mg per dose under the skin every 4 weeks Inject 300mg per dose under the skin every 4 weeks Inject 450mg per dose under the skin every 4 weeks Inject 600mg per dose under the skin every 4 weeks Inject other: _____ mg per dose under the skin every 4 weeks Every 2 weeks dosing: Inject 225mg per dose under the skin every 2 weeks Inject 300mg per dose under the skin every 2 weeks Inject 375mg per dose under the skin every 2 weeks Inject 450mg per dose under the skin every 2 weeks Inject 525mg per dose under the skin every 2 weeks Inject 600mg per dose under the skin every 2 weeks Inject other: _____ mg per dose under the skin every 2 weeks	1-month supply 3-month supply Other: _____ Refills _____
Epinephrine/EpiPen®	0.3mg IM as needed for anaphylaxis 0.15mg IM as needed for anaphylaxis		1-month supply Refill x 1 year unless noted otherwise Other: _____
Xolair vial supplies: Sterile water for injection 10mL vial for reconstitution QS per doses Administration Supply Kit consisting of: <ul style="list-style-type: none"> • Alcohol swabs • Flexible bandages 1" x 3" • 3mL Luer Lock injection syringe • NDL 18G x 1 1/2" Safety Glide needle for reconstitution • NDL 25G x 5/8" Safety Glide needle for subcutaneous injection No supplies (Supplies will be sent with shipment unless indicated.)			Send quantity sufficient for medication days supply

Xolair Self-Administration Physician authorization to ship to the home

Has the patient received at least 3 doses of Xolair under the guidance of a healthcare provider without hypersensitivity reactions and the healthcare provider has completed the assessment of risk for anaphylaxis and mitigation strategies, and has determined that self-administration is appropriate? Yes No

If shipped to physician's office or infusion clinic, physician accepts on behalf of patient for administration in office or infusion clinic.

Prescriber's signature required (sign below) (Physician attests this is his/her legal signature. NO STAMPS)

**SIGN
HERE**

Date

Dispense as written

Date

Substitution allowed

The prescriber is to comply with his/her state-specific prescription requirements such as e-prescribing, state-specific prescription form, fax language, etc. Non-compliance with state-specific requirements could result in outreach to the prescriber.