

# Rheumatoid arthritis—Intravenous

Four simple steps to submit your referral.

## 1 PATIENT INFORMATION

New patient  Current

Patient's name \_\_\_\_\_  
 Date of birth \_\_\_\_\_  Male  Female Last 4 digits of SSN \_\_\_\_\_  
 Street address \_\_\_\_\_ Apt # \_\_\_\_\_  
 City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_  
 Parent/guardian (if applicable) \_\_\_\_\_  
 Home phone \_\_\_\_\_ Work phone \_\_\_\_\_ Cell phone \_\_\_\_\_  
 Evening phone \_\_\_\_\_ E-mail address \_\_\_\_\_  
 Patient's primary language:  English  Other If other, please specify \_\_\_\_\_

Please attach front and back of patient's insurance cards or complete information below.

Insurance company \_\_\_\_\_ Phone \_\_\_\_\_  
 Insured's name \_\_\_\_\_  
 Insured's employer \_\_\_\_\_ Relationship to patient \_\_\_\_\_  
 Identification # \_\_\_\_\_ Policy/group # \_\_\_\_\_  
 Prescription card:  Yes  No If yes, carrier \_\_\_\_\_  
 Policy # \_\_\_\_\_ Group # \_\_\_\_\_ Is patient eligible for Medicare?  Yes  No  
 Does patient have a secondary insurance?  Yes  No

## 2 PRESCRIBER INFORMATION

All fields must be completed to expedite prescription fulfillment.

Date \_\_\_\_\_ Time \_\_\_\_\_ Date medication needed \_\_\_\_\_  
 Prescriber's name and title \_\_\_\_\_  
 If NP or PA, under direction of Dr. \_\_\_\_\_  
 Office contact and title \_\_\_\_\_  
 Clinic/hospital affiliation \_\_\_\_\_ Street address \_\_\_\_\_ Suite # \_\_\_\_\_  
 City \_\_\_\_\_ State \_\_\_\_\_ Zip \_\_\_\_\_ Phone \_\_\_\_\_ Fax \_\_\_\_\_  
 NPI # \_\_\_\_\_ License # \_\_\_\_\_  
 Deliver product to:  Office  Patient's home  Clinic  
 Clinic location \_\_\_\_\_

## 3 CLINICAL INFORMATION

Primary ICD-10 code: \_\_\_\_\_  
 Has the patient been treated previously for this condition?  Yes  No  
 Is patient currently on therapy?  Yes  No  
 Please list all therapies tried/failed: \_\_\_\_\_  
 \_\_\_\_\_  
 Patient wt \_\_\_\_\_  Date wt obtained \_\_\_\_\_  
 NKDA  Known drug allergies \_\_\_\_\_  
 \_\_\_\_\_  
 Concurrent meds \_\_\_\_\_

Please fax completed form to the RA team at 888.302.1028.

To reach your team, call toll-free 888.608.9010.

The document(s) accompanying this transmission may contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender immediately and arrange for the return or destruction of these documents.

All rights in the product names, trade names or logos of all third-party products that appear in this form, whether or not appearing with the trademark symbol, belong exclusively to their respective owners.

© 2018 Accredo Health Group, Inc. An Express Scripts Company. All Rights Reserved.  
 RAS-00027-012418 amc9312 CRP17A2\_0618

## 4 PRESCRIBING INFORMATION

Medication	Strength / Formulation	Directions	Quantity/Refills
<input type="checkbox"/> Actemra® (tocilizumab)	<input type="checkbox"/> 4 mg/kg intravenous infusion every 4 weeks. Maximum dose of 800 mg/infusion <input type="checkbox"/> 8 mg/kg intravenous infusion every 4 weeks. Maximum dose of 800 mg/infusion	Dilute desired dose with normal saline to a total volume of 100 mL to be infused over 1 hour.	<input type="checkbox"/> 1-month supply <input type="checkbox"/> 3-month supply <input type="checkbox"/> Other _____ Refills _____
<input type="checkbox"/> Orencia® (abatacept)	<input type="checkbox"/> 500 mg (less than 60 kg) <input type="checkbox"/> 750 mg (60–100 kg) <input type="checkbox"/> 1000 mg (over 100 kg) <input type="checkbox"/> Juvenile arthritis 10 mg/kg if less than 75 kg Starting dose: <input type="checkbox"/> at week: 0, 2 and 6, then every 4 weeks Maintenance dose: <input type="checkbox"/> every 4 weeks	Reconstitute each vial of Orencia with 10 mL of sterile water. Dilute desired dose to total of 100 mL in normal saline to be infused over 30 minutes.	<input type="checkbox"/> 1-month supply <input type="checkbox"/> 3-month supply <input type="checkbox"/> Other _____ Refills _____
<input type="checkbox"/> Remicade® (infliximab)	Starting dose: <input type="checkbox"/> 5 mg/kg _____ mg IV at week: 0, 2, 6 <input type="checkbox"/> 3 mg/kg _____ mg IV at week: 0, 2, 6 <input type="checkbox"/> Other _____ Maintenance dose: (_____ mg/kg) _____ mg IV every _____ weeks	Reconstitute each vial of Remicade with 10 mL of sterile water. Dilute desired dose to total of 250 mL in normal saline to be infused over a period NOT less than 2 hours.	<input type="checkbox"/> 1-month supply <input type="checkbox"/> 3-month supply <input type="checkbox"/> Other _____ Refills _____
<input type="checkbox"/> Simponi Aria® (golimumab)	Starting dose: <input type="checkbox"/> 2 mg/kg _____ mg IV at week: 0, 4 and every 8 weeks <input type="checkbox"/> Other _____ Maintenance dose: <input type="checkbox"/> 2 mg/kg _____ mg IV every 8 weeks <input type="checkbox"/> Other _____	Each 4 mL vial contains 50 mg of Simponi Aria. Dilute the total desired volume (based on 2 mg/kg dosing) of Simponi Aria with 0.9% sodium chloride to a total volume of 100 mL. Infuse diluted solution over a period of 30 minutes.	<input type="checkbox"/> 1-month supply <input type="checkbox"/> 3-month supply <input type="checkbox"/> Other _____ Refills _____
<input type="checkbox"/> Other _____			<input type="checkbox"/> 1-month supply <input type="checkbox"/> 3-month supply <input type="checkbox"/> Other _____ Refills _____

Complete the below information if assistance from Accredo is requested in the coordination of your patient's infusion therapy.

Preferred infusion setting:  Home  Infusion clinic

**Premedication orders**  
 Acetaminophen 650 mg PO 30 min prior to infusion  
 Diphenhydramine 50 mg PO 30 min prior to infusion  
 Hydrocortisone 100 mg IV PO 30 min prior to infusion  
 Other \_\_\_\_\_

**Hypersensitivity/anaphylaxis orders**  Stop infusion  Start normal saline at TKO

**Medicate with:**  
 Epinephrine/EpiPen® 0.3 mg IM as needed for anaphylaxis.  
 Diphenhydramine 50 mg slow IVP PRN for urticaria, pruritis, SOB. Administer IM if there is no IV access.  
 Hydrocortisone 100 mg slow IVP PRN for urticaria, pruritis, SOB. Administer IM in there is no IV access.  
 Solumedrol 125 mg slow IVP PRN for urticaria, pruritis, SOB. Administer IM if there is no IV access.

**For anaphylactic reaction, activate 911.** Notify physician of type reaction and action taken. Verbal report and transfer care to EMS, if applicable.

**Flushing orders**  
 Peripheral access  
 Central venous access  
 0.9% sodium chloride flush with \_\_\_\_\_ mL IV before and after medication and IVP for maintenance.  
 Heparin \_\_\_\_\_ units per mL. Flush with \_\_\_\_\_ units as final flush and as directed.

**Lab orders**  
 Skilled nursing visit as needed to establish venous access, administer medication and assess general status and response to therapy. Dispense 1 month of drug, flushes, needles, syringes, ancillary supplies and medical equipment necessary to establish access and administer medication.

Prescription to include all necessary ancillary supplies (needles, syringes, etc.)  
 If shipped to physician's office, physician accepts on behalf of patient for administration in office.

By signing below, I certify that the above therapy is medically necessary.  
 Prescriber's signature (sign below) (Prescriber attests this is his/her legal signature. **NO STAMPS**)

Date \_\_\_\_\_ Substitution allowed \_\_\_\_\_ Date \_\_\_\_\_ Dispense as written \_\_\_\_\_

The prescriber is to comply with his/her state-specific prescription requirements such as e-prescribing, state-specific prescription form, fax language, etc. Non-compliance with state-specific requirements could result in outreach to the prescriber.

PHYSICIAN SIGNATURE REQUIRED